

## Louisiana Tech University Louisiana Tech Digital Commons

---

ANS Research Symposium

ANS Research Symposium 2019

---

Apr 11th, 8:30 AM - 11:30 AM

# Determining Expression Levels of the Notch Signaling Pathway in Self-Renewing hASCs

John B. Cart  
*Louisiana Tech University*

Avery Bryan  
*Louisiana Tech University*

Chris Miller  
*Louisiana Tech University*

Jamie Newman  
*Louisiana Tech University*

Follow this and additional works at: <https://digitalcommons.latech.edu/ans-research-symposium>

---

### Recommended Citation

Cart, John B.; Bryan, Avery; Miller, Chris; and Newman, Jamie, "Determining Expression Levels of the Notch Signaling Pathway in Self-Renewing hASCs" (2019). *ANS Research Symposium*. 32.  
<https://digitalcommons.latech.edu/ans-research-symposium/2019/poster-presentations/32>

This Event is brought to you for free and open access by the Conferences and Symposia at Louisiana Tech Digital Commons. It has been accepted for inclusion in ANS Research Symposium by an authorized administrator of Louisiana Tech Digital Commons. For more information, please contact [digitalcommons@latech.edu](mailto:digitalcommons@latech.edu).

# Determining Expression Levels of the Notch Signaling Pathway in Self-Renewing hASCs

John Bradley Cart<sup>1</sup>, Avery Bryan<sup>1</sup>, Chris Miller<sup>2</sup>, Dr. Jamie Newman<sup>3</sup>

<sup>1</sup>*Undergraduate Biology, Louisiana Tech University*

<sup>2</sup>*Graduate Student, Molecular Science and Nanotechnology, Louisiana Tech University*

<sup>3</sup>*Assistant Professor, School of Biological Sciences*

---

Human adipose-derived stem cells (hASCs) have immense potential for regenerative medicine. They can potentially differentiate to form bone, muscle, or cartilage tissue making them very promising in combating degenerative diseases. One of the key elements for maximizing the clinical usage of these cells is determining mechanisms that determine cell fate. The Notch signaling pathway is highly conserved across multiple species and is made up of 4 receptors (Notch 1,2,3, and 4) and 5 ligands (Jag1, Jag2, Dll1, Dll3, and Dll4) each performing an important role in the process of cellular homeostasis and differentiation. Although this pathway is known to be significant in hASC differentiation, very little is known about the role of each receptor and how they work together to maintain and direct cell state. If hASCs are to be used in a clinical setting, then the Notch pathway must be fully understood. We investigate this pathway by determining the expression levels of each of its receptors and ligands. This allows us to piece together how the pathway works by determining how involved these proteins are in the Notch pathway. We further investigate the mechanism of this pathway by performing an siRNA mediated knock-down of Notch 3, a protein which has been determined to play a key role in the pathway. The siRNA will interrupt the translation of the Notch 3 gene from mRNA to protein which will lead to reduced expression of Notch 3. We will then look at the expression of other components of the Notch pathway. If there is a change in expression level in a protein after the Notch 3 knock-down, then there is likely a connection between that protein and Notch 3. By piecing together all of this information, we can begin to get a complete picture of the mechanisms driving the Notch signaling pathway. Once this pathway is understood, it could lead to the development of treatments for degenerative diseases using hASCs.